

30.8.2024

Näytteenotto ulosteen bakteeripatogeenien nukleiinihappo-osoitusta varten (F-BaktNhO, 54328, F-MikrNhO 54341) ENGLANTI

Stool sample for measuring nucleic acid in pathogenic bacteria (F –BaktNhO 54328, F-MikrNhO 54341)

Sampling kit

- Bacteria transportation tube
- Stool specimen container
- Labels for personal data and travel information

Sample collection

- Time the collection so that you can take the sample to the laboratory on the same day.
- Empty your bladder (there should not be urine in the specimen).
- Pass stool on toilet paper or into a clean, dry dish, for example a disposable dish or a potty or other suitable dish covered with a clean plastic bag (e.g. freezer bag).
- Transfer stool both into the bacteria transportation tube and the stool specimen container according to the following instructions:

Bacteria transportation tube

- Remove the bacteria transportation tube from the plastic cover.
- Remove the plastic propective plug from top of the tube (plug can be thrown to waste).
- Dip the cotton head of the sample stick well into the stool, preferable in the slimy or bloody part.
- Insert the sample stick into the bacterial transportation tube so that the cotton head immerses into the gel and the plug at the end of the stick closes the tube tightly.

Stool specimen container

- Put fresh stool into the sample container (about half the container).
- Screw the top of the specimen container shut tightly and make sure the container does not leak.
- Write your name, personal identity code and date and time of sample collection as well as information about your travel abroad on the labels:
 - If you have travelled abroad in the past few weeks, write: COUNTRY and RETURN DATE.
 - If you have not travelled abroad in the past few weeks, write: NO TRAVELS.
- Stick the labels lengthwise both on the bacteria transportation tube and the stool specimen container. Put both in a plastic bag. If there is a paper referral, leave it outside the bag

Storage

Keep the sample waiting to be transported in a cold place (refrigerator). Take the tube and potential paper referral to the laboratory on the day of sample collection in room temperature or in refrigerator temperature.